

AN OVERVIEW OF VARIABLE EXPONENT LEBESGUE AND SOBOLEV SPACES

PETTERI HARJULEHTO AND PETER HÄSTÖ

ABSTRACT. In this paper we give an overview of the development of variable exponent Lebesgue and Sobolev spaces from the beginning of the 1990s till today. We describe the basic theory and mention some applications. Additionally, we prove a new result on the density of smooth functions in variable exponent Sobolev spaces on the real line.

1. INTRODUCTION

In recent years there has been increasing interest toward variable exponent Lebesgue and Sobolev spaces. Although similar spaces had been previously considered by some investigators (e.g., H. Hudzik [33–35] and J. Museliak [43]) the present line of investigation goes back to a paper by O. Kováčik and J. Rákosník [40] from 1991. After this paper not much happened until the late 1990s. At this point the subject seems to have been independently (re)discovered by several researchers: S. Samko [49–51] working based on earlier Russian work (I. Sharapudinov [52] and V. Zhikov [53]), X. Fan and collaborators [16–22] and G. Mingione and collaborators [1–4, 42] drawing inspiration from the study of differential equations (e.g. Marcellini [41]). Around the turn of the millennium also researchers basing their efforts directly on the work of Kováčik and Rákosník got new momentum with the investigations of D. Edmunds and collaborators [10–14] L. Diening [6, 7] and others [5, 8, 9, 15, 44, 47]. Our group at the University of Helsinki started its research in the Fall of 2002. To date our efforts have yielded some publications [25, 28] and preprints [26, 27, 29–31] The last couple of years have seen a (partial) integration of the separate lines of investigation, but much still remains to be done. We believe that variable exponent spaces are a good entry-point requiring little prerequisites for someone who wants to start research on Sobolev spaces.

The main incentive for many of the investigators of variable exponent spaces is relaxing the coercivity conditions assumed for the solutions of a differential equation or the corresponding variational integral. Typically one would assume that the function $F: G \times \mathbb{R}^n \rightarrow [0, \infty)$ in the integral

$$(1.1) \quad \int_G F(x, \nabla u(x)) dx$$

satisfies the coercivity condition

$$(1.2) \quad \frac{1}{c} |\nabla u(x)|^p \leq F(x, \nabla u(x)) \leq c |\nabla u(x)|^p.$$

In this case the minimizers of (1.1) are in $W^{1,p}(G)$, [32]. (We ignore many technical assumptions here, see the reference for details.) Now if we just replace p by $p(x)$ in (1.2), then we get a more general condition, but in what space will the minimizers of the variational

Date: September 26, 2003.

2000 Mathematics Subject Classification. 46E35.

Key words and phrases. Variable exponent, Lebesgue space, Sobolev space, density of smooth functions.

integral lie now? Although the theory is not yet complete the tenet of variable-exponent-space investigators is that it will be $W^{1,p(\cdot)}(G)$, the variable exponent Sobolev space (defined below).

Perhaps the next question that comes to mind is: why should we want to generalize (1.2) in the first place? The answer is again somewhat tentative – we believe that it will be useful in modeling non-homogeneous material. Until now, only one such application has been investigated in greater detail, electrorheological fluids. These fluids have the interesting property that their viscosity depends on the electric field in the fluid. For a general account of the underlying physics confer [24] and for some technical applications [45]. The mathematical theory was presented in the monograph [48] by M. Růžička, see also [3].

The remaining part of this article strives to give a little more detailed account of the mathematics of variable exponent spaces. In the next section we present the theory of Lebesgue spaces, in Section 3 that of Sobolev spaces. In the final section we present the proof of a new result on the density of smooth functions in Sobolev spaces in the one-dimensional case. This problem is a good example of how something which is easy in classical Sobolev spaces turns out to be quite difficult in the variable case. The proof also illustrates some techniques that can be used in variable exponent spaces.

2. LEBESGUE SPACES

Let $p : \mathbb{R}^n \rightarrow [1, \infty)$ be a measurable function (called the *variable exponent* on \mathbb{R}^n). Throughout this paper the function p always denotes a variable exponent; also, we define $p^+ = \operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x)$ and $p^- = \operatorname{ess\,inf}_{x \in \mathbb{R}^n} p(x)$. We define the *variable exponent Lebesgue space* $L^{p(\cdot)}(\mathbb{R}^n)$ to consist of all measurable functions $u : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$\varrho_{p(\cdot)}(\lambda u) = \int_{\mathbb{R}^n} |\lambda u(x)|^{p(x)} dx < \infty$$

for some $\lambda > 0$. The function $\varrho_{p(\cdot)} : L^{p(\cdot)}(\mathbb{R}^n) \rightarrow [0, \infty)$ is called the *modular* of the space $L^{p(\cdot)}(\mathbb{R}^n)$. We define a norm, the so-called *Luxemburg norm*, on this space by the formula

$$\|u\|_{p(\cdot)} = \inf\{\lambda > 0 : \varrho_{p(\cdot)}(u/\lambda) \leq 1\}.$$

If p is a constant function, then the variable exponent Lebesgue spaces coincides with the classical Lebesgue space and so the notation can give rise to no confusion. The variable exponent Lebesgue spaces is a special case of Orlicz–Musielak spaces treated by Musielak [43]. The basic properties of these spaces can be found in the paper of Kováčik and Rákosník [40]; many of these properties were independently established by Fan and Zhao [21].

2.1. Basic properties. Variable exponent Lebesgue spaces resemble classical Lebesgue spaces in many respects – they are Banach spaces [40, Theorem 2.5], the Hölder inequality holds [40, Theorem 2.1], they are reflexive if and only if $1 < p^- \leq p^+ < \infty$ [40, Corollary 2.7] and continuous functions are dense if $p^+ < \infty$ [40, Theorem 2.11]. The inclusion between Lebesgue spaces also generalizes naturally [40, Theorem 2.8]: if $0 < |\Omega| < \infty$ and p, q are variable exponent so that $p(x) \leq q(x)$ almost everywhere in Ω then there exists an imbedding $L^{q(\cdot)}(\Omega) \hookrightarrow L^{p(\cdot)}(\Omega)$ whose norm does not exceed $|\Omega| + 1$.

One problem in variable exponent Lebesgue spaces is the relation between the modular and the norm. If $p^+ < \infty$ then

$$\min\{\|u\|_{p(\cdot)}^{p^-}, \|u\|_{p(\cdot)}^{p^+}\} \leq \varrho_{p(\cdot)}(u) \leq \max\{\|u\|_{p(\cdot)}^{p^-}, \|u\|_{p(\cdot)}^{p^+}\}$$

[21, Theorem 1.3]. It follows that if $p^+ < \infty$ and (f_i) is a sequence of functions in $L^{p(\cdot)}$, then $\|f_i\|_{p(\cdot)} \rightarrow 0$ if and only if $\varrho_{p(\cdot)}(f_i) \rightarrow 0$ [40, Theorem 2.4]. However, if $p^+ = \infty$ the topology defined by the norm is not the same as that defined by the modular. Spaces with $p^+ = \infty$ have been studied by Edmunds, Lang and Nekvinda [10].

Variable exponent Lebesgue spaces do not have the mean continuity property. If p is a non-constant continuous function in an open ball B , then there exists a function $u \in L^{p(\cdot)}(B)$ such that $u(x+h) \notin L^{p(\cdot)}(B)$ for some $h \in \mathbb{R}^n$ with arbitrary small norm [40, Theorem 2.10]. In [23], Fiorenza has studied a weak version of mean continuity in variable exponent Sobolev spaces.

The Hardy–Littlewood maximal operator. Finally we give an example of a classical property which is now understood quite well also in the variable exponent case. Assume that $1 < p^- \leq p^+ < \infty$ and there exists a constant $C > 0$ such that

$$(2.1) \quad |p(x) - p(y)| \leq \frac{C}{-\log|x-y|}$$

for every $x, y \in \mathbb{R}^n$ with $|x - y| \leq \frac{1}{2}$, and

$$|p(x) - p(y)| \leq \frac{C}{\log(e+|x|)}$$

for every $x, y \in \mathbb{R}^n$ with $|y| \geq |x|$. Under these assumptions on p , Cruz-Uribe, Fiorenza and Neugebauer proved that the Hardy–Littlewood maximal operator is bounded from $L^{p(\cdot)}(\mathbb{R}^n)$ to itself [5, Theorem 1.5]. This was an improvement of earlier work by Diening [6] and Nekvinda [44]. The assumptions on p are more or less sharp: Pick and Růžička showed that if (2.1) does not hold then the Hardy–Littlewood maximal operator is not bounded [46] while Cruz-Uribe, Fiorenza and Neugebauer proved that if p is upper semicontinuous and $p^- = 1$ then the operator is not bounded [5, Theorem 1.7]. Cruz-Uribe, Fiorenza and Neugebauer proved a weak type estimate of the Hardy–Littlewood maximal operator provided that $1/p \in RH_\infty$ [5, Theorem 1.8]. This assumption is true for example if $p^+ < \infty$. Maximal operators have also been studied in weighted $L^{p(\cdot)}$ spaces by Kokilashvili and Samko [39] and in metric measure spaces by Harjulehto, Hästö and Pere [31].

Samko [49, 50], Kokilashvili and Samko [36, 37, 38] and Diening and Růžička [8, 9] have also studied other kinds of integral operators in variable exponent and weighted variable exponent spaces.

3. SOBOLEV SPACES

The *variable exponent Sobolev space* $W^{1,p(\cdot)}(\mathbb{R}^n)$ is the subspace of functions $L^{p(\cdot)}(\mathbb{R}^n)$ whose distributional gradient exists almost everywhere and lies in $L^{p(\cdot)}(\mathbb{R}^n)$. The function $\varrho_{1,p(\cdot)} : W^{1,p(\cdot)}(\mathbb{R}^n) \rightarrow [0, \infty)$ is defined by $\varrho_{1,p(\cdot)}(u) = \varrho_{p(\cdot)}(u) + \varrho_{p(\cdot)}(|\nabla u|)$. The norm $\|u\|_{1,p(\cdot)} = \|u\|_{p(\cdot)} + \|\nabla u\|_{p(\cdot)}$ makes $W^{1,p(\cdot)}(\mathbb{R}^n)$ a Banach space, separable if $p^+ < \infty$ and reflexive if $1 < p^- \leq p^+ < \infty$ [40, Theorem 3.1]. Variable exponent Sobolev spaces have close connection to the study of partial differential equations and variational integrals with non-standard growth, see for example [1, 2, 4, 20, 53]

Density of smooth functions. Samko [51] and Fan and Zhao [20] have proven independently that if p satisfies (2.1) then $C_0^\infty(\mathbb{R}^n)$ is dense in $W^{1,p(\cdot)}(\mathbb{R}^n)$. Edmunds and Rákosník showed that if $\Omega \subset \mathbb{R}^n$ is bounded with Lipschitz boundary and the exponent p satisfies a condition similar to (2.1) then $C^\infty(\bar{\Omega})$ is dense in $W^{1,p(\cdot)}(\Omega)$ [13, Theorem 3.2], see also Diening’s version of this result [6, Theorem 3.7].

Edmunds and Rákosník derived the same conclusion under a local monotony condition on p [12, Theorem 1]. In Section 4 of this paper we show that on the real line C_0^∞ -functions are dense in $W^{1,p(\cdot)}(\mathbb{R})$ provided only that $p^+ < \infty$. An example constructed by Zhikov [53, p. 36] shows that if the exponent is discontinuous then smooth functions need not be dense in variable exponent Sobolev space if the dimension is at least 2.

The Sobolev and Poincaré inequalities. Diening proved that if $1 < p^- \leq p^+ < n$, (2.1) holds and p is constant outside some large ball then there exists a continuous imbedding $W^{1,p(\cdot)}(\mathbb{R}^n) \hookrightarrow L^{p^*(\cdot)}(\mathbb{R}^n)$ [7, Theorem 5.2]. Here $p^*(x) = np(x)/(n - p(x))$. He also has a version of this in bounded domains with Lipschitz boundary [7, Corollary 5.3]. Fan, Shen and Zhao have proven the Sobolev imbedding for a larger class of domains than Diening but under stricter condition on p [16]. Edmunds and Rákosník assumed that p is Lipschitz continuous in the closure of an open set Ω and there exists real numbers q, r such that $1 < r \leq p(x) \leq q < \min\{n, r^*\}$ and showed

$$\|u\|_{p^*(\cdot)} \leq C \|\nabla u\|_{p(\cdot)}$$

for every $u \in W^{1,p(\cdot)}(\Omega)$ with $\text{spt } u \subset \Omega$ [13, Theorem 3.1]. Kováčik and Rákosník gave an example which shows that the Sobolev imbedding does not hold if the exponent is discontinuous [40, Example 3.2].

The situation changes if we are ready to sacrifice an $\varepsilon > 0$ from one of the norms involved. Kováčik and Rákosník [40, Theorem 3.3] assumed that $p : \bar{\Omega} \rightarrow [1, n]$ is continuous and showed that for every small $\varepsilon > 0$ and every measurable function $1 \leq q(x) \leq p^*(x) - \varepsilon$ the inequality

$$\|u\|_{q(\cdot)} \leq C \|\nabla u\|_{p(\cdot)}, \quad u \in C_0^\infty(\Omega)$$

holds. Edmunds and Rakosnik also have Sobolev type inequalities in variable exponent spaces with weights [14]. Sobolev inequalities are discussed at length in [47]. The Sobolev imbedding for $p(x) > n$ has been studied in [25, Sections 3 and 4] by Harjulehto and Hästö.

It is interesting to note that the Poincaré inequality holds under a much weaker assumption on p than the Sobolev inequality and imbedding, namely if the exponent is not too discontinuous [25, Section 2]. For instance, if the exponent is continuous up to the boundary in a bounded convex domain, then we have a Poincaré inequality [25, Corollary 2.16]. Harjulehto and Hästö also gave an example which shows that the condition for p is sharp [25, Example 2.6].

Sobolev capacity and its applications. In [28] Harjulehto, Hästö, Koskenoja and Varonen introduced a Sobolev capacity in the variable exponent Sobolev space, which is defined as follows. Suppose that E is an arbitrary subset of \mathbb{R}^n . The *Sobolev $p(\cdot)$ -capacity* of E is defined by

$$C_{p(\cdot)}(E) = \inf \int_{\mathbb{R}^n} (|u(x)|^{p(x)} + |\nabla u(x)|^{p(x)}) dx,$$

where the infimum is taken over those $u \in W^{1,p(\cdot)}(\mathbb{R}^n)$ which are at least one in some open set containing E . If $1 < p^- \leq p^+ < \infty$, then the Sobolev $p(\cdot)$ -capacity is an outer measure and Choquet capacity [28, Corollaries 3.3 and 3.4]. It is possible to relax the conditions on p by assuming $p^- > 1$ locally [30]. As in the fixed exponent case the capacity is a finer measure than the n -dimensional Lebesgue measure [28, Section 4]. If continuous functions are dense in variable exponent Sobolev spaces and if $1 < p^- \leq p^+ < \infty$ then every Sobolev function has a quasicontinuous representative [28, Theorem 5.2].

Using the Sobolev $p(\cdot)$ -capacity Harjulehto, Hästö, Koskenoja and Varonen defined Sobolev spaces with zero boundary values as in metric measure spaces. This definition coincides with that based on the closure of C_0^∞ -functions if the smooth functions are dense [28, Chapter 3]. This allowed them to study the existence of a minimizer of the Dirichlet energy integral in variable exponent spaces with Sobolev boundary values [29]. This problem is not trivial even in the one-dimensional case, see [27]. The Sobolev $p(\cdot)$ -capacity also seems to be the right one for studying Lebesgue points of Sobolev functions. Harjulehto and Hästö proved in [26] that quasiaevery point is a Lebesgue point of a Sobolev function if p satisfies (2.1). It is also possible to define a capacity using the norm instead of the modular [25, Theorem 4.2] however this does not yield a Choquet capacity.

Summary. We have seen that condition (2.1) is a very common assumption in the theory of variable exponent spaces. This condition, sometimes strengthened with some technical assumptions, guaranties that many classical properties hold. Moreover, it seems that in many instances this is exactly the right condition. However, there are also many cases in which there is a gap between the necessary and sufficient conditions that we cannot bridge.

Under the weaker assumptions $p^+ < \infty$ or $1 < p^- \leq p^+ < \infty$ variable exponent spaces still have many interesting properties. However, in this case many classical techniques (e.g. convolution) are not available and new methods must be devised to prove standard results. For $p^+ = \infty$ only a few properties have been established for these spaces.

4. DENSITY OF SMOOTH FUNCTIONS ON THE REAL LINE

In this section we show that the density problem, which remains open in higher dimensions despite much effort, is essentially trivial in the one-dimensional case. As for the condition $p^+ < \infty$, note that for $p^+ = \infty$ we do not even know if $C^\infty(I) \cap L^{p(\cdot)}(I)$ is dense in $L^{p(\cdot)}(I)$. Here I is the open interval $(-1, 1)$.

4.1. Theorem. *If $p^+ < \infty$, then $C^\infty(I) \cap W^{1,p(\cdot)}(I)$ is dense in $W^{1,p(\cdot)}(I)$.*

Proof. Let $u \in W^{1,p(\cdot)}(I)$ and fix $\epsilon > 0$. By [40, Theorem 3.4] there exists a function $g \in C^\infty(I) \cap L^{p(\cdot)}(I)$ such that $\|u' - g\|_{p(\cdot)} \leq \epsilon$. Fix $z \in I$. Define

$$v(y) = u(z) + \int_z^y g(x) dx$$

for $y \in I$, so that $v' = g$. We then have

$$|v(y) - u(y)| \leq \left| u(z) + \int_z^y g(x) dx - u(z) - \int_z^y u'(x) dx \right| \leq \int_z^y |g(x) - u'(x)| dx \leq \|u' - g\|_1.$$

By [40, Theorem 2.8] it follows that $\|u' - g\|_1 \leq 2\|u' - g\|_{p(\cdot)} < 2\epsilon$. By the same theorem and the above inequality, $\|v - u\|_{p(\cdot)} \leq 2\|v - u\|_\infty < 4\epsilon$, and so

$$\|v - u\|_{1,p(\cdot)} = \|v - u\|_{p(\cdot)} + \|u' - g\|_{p(\cdot)} < 5\epsilon.$$

Since $v \in C^\infty(I)$ and since ϵ can be arbitrarily small, we have found our C^∞ approximation and so we are done. \square

4.2. Theorem. *If $p^+ < \infty$, then $C_0^\infty(\mathbb{R})$ is dense in $W^{1,p(\cdot)}(\mathbb{R})$.*

Proof. Let $u \in W^{1,p(\cdot)}(\mathbb{R})$ and fix $\epsilon > 0$. We denote by I_x the open interval $(x - 1, x + 1)$ for $x \in \mathbb{R}$. Let $\phi(x) \in C_0^\infty([-1, 1])$ (extended by 0 outside $[-1, 1]$) be a non-negative L -Lipschitz function such that $\sum_{k \in \mathbb{Z}} \phi(x - k)$ is identically equal to 1.

For $k \in \mathbb{Z}$ we choose a $C^\infty(I_k)$ function v_k such that $\|u - v_k\|_{W^{1,p(\cdot)}(I_k)} < \epsilon/2^{|k|}$, using Theorem 4.1. Define $v(x) = \sum_{k \in \mathbb{Z}} v_k(x)\phi(x - k)$. Then

$$\begin{aligned} \|u - v\|_{1,p(\cdot)} &= \left\| \sum_{k \in \mathbb{Z}} (u(x) - v_k(x))\phi(x - k) \right\|_{1,p(x)} \\ &\leq \sum_{k \in \mathbb{Z}} \|(u(x) - v_k(x))\phi(x - k)\|_{1,p(x)} \\ &\leq \sum_{k \in \mathbb{Z}} \|u - v_k\|_{p(\cdot)} + \left\| \{(u(x) - v_k(x))\phi(x - k)\}' \right\|_{p(x)} \\ &\leq \sum_{k \in \mathbb{Z}} (1 + L)\|u - v_k\|_{p(\cdot)} + \|(u - v_k)'\|_{p(\cdot)} \\ &\leq (1 + L) \sum_{k \in \mathbb{Z}} \|u - v_k\|_{1,p(\cdot)} \\ &\leq (1 + L)\epsilon \sum_{k \in \mathbb{Z}} 2^{-|k|} = 3(1 + L)\epsilon. \end{aligned}$$

Since

$$\lim_{l \rightarrow \infty} \left\| u(x) - \sum_{k=-l}^l v_k(x)\phi(x - k) \right\|_{1,p(x)} \leq 3(1 + L)\epsilon,$$

it follows that there exists an integer l such that

$$\left\| u(x) - \sum_{k=-l}^l v_k(x)\phi(x - k) \right\|_{1,p(x)} \leq 4(1 + L)\epsilon.$$

But clearly $\sum_{k=-l}^l v_k(x)\phi(x - k) \in C_0^\infty(\mathbb{R})$, so we are done. \square

Acknowledgement

The authors wish to thank the referee for a very meticulous reading of the manuscript.

REFERENCES

- [1] E. Acerbi and G. Mingione: Regularity results for a class of functionals with non-standard growth, *Arch. Ration. Mech. Anal.* **156** (2001), 121–140.
- [2] E. Acerbi and G. Mingione: Regularity results for stationary electro-rheological fluids, *Arch. Ration. Mech. Anal.* **164** (2002), 213–259.
- [3] E. Acerbi and G. Mingione: Regularity results for electrorheological fluids: the stationary case *C. R. Acad. Sci. Paris* **334** (2002), 817–822.
- [4] A. Coscia and G. Mingione: Hölder continuity of the gradient of $p(x)$ -harmonic mappings, *C. R. Acad. Sci. Paris*, **328** (1999), 363–368.
- [5] D. Cruz-Uribe, A. Fiorenza and C. J. Neugebauer: The maximal function on variable L^p spaces, *Ann. Acad. Sci. Fenn.* **28** (2003), 223–238.
- [6] L. Diening: Maximal function on generalized Lebesgue spaces $L^{p(\cdot)}$, *Math. Inequal. Appl.*, to appear.
- [7] L. Diening: Riesz Potential and Sobolev Embeddings of generalized Lebesgue and Sobolev Spaces $L^{p(\cdot)}$ and $W^{k,p(\cdot)}$, *Math. Nachr.*, to appear.
- [8] L. Diening and M. Růžička: Calderón-Zygmund operators on generalized Lebesgue spaces $L^{p(\cdot)}$ and problems related to fluid dynamics, *J. Reine Ang. Math.*, to appear.

- [9] L. Diening and M. Růžička: Integral operators on the halfspace in generalized Lebesgue spaces $L^{p(\cdot)}$, preprint (2003).
- [10] D. E. Edmunds, J. Lang and A. Nekvinda: On $L^{p(x)}$ norms, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* **455** (1999), no. 1981, 219–225.
- [11] D. E. Edmunds and A. Meskhi: Potential-type operators in $L^{p(x)}$ spaces, *Z. Anal. Anwendungen* **21** (2002), no. 3, 681–690.
- [12] D. E. Edmunds and J. Rákosník: Density of smooth functions in $W^{k,p(x)}(\Omega)$, *Proc. Roy. Soc. London Ser. A* **437** (1992), 229–236.
- [13] D. E. Edmunds and J. Rákosník: Sobolev embedding with variable exponent, *Studia Math.* **143** (2000), 267–293.
- [14] D. E. Edmunds and J. Rákosník: Sobolev embedding with variable exponent, II, *Math. Nachr.* **246-247** (2002), 53–67.
- [15] M. Eleuteri: Hölder continuity results for a class of functionals with non standard growth, *Boll. Unione Mat. Ital.*, to appear.
- [16] X. Fan, J. Shen and D. Zhao: Sobolev embedding theorems for spaces $W^{k,p(x)}$, *J. Math. Anal. Appl.* **262** (2001), 749–760.
- [17] X.-L. Fan, H.-Q. Wu and F.-Z. Wang: Hartman-type results for $p(t)$ -Laplacian systems, *Nonlinear Anal.* **52** (2003), 585–594.
- [18] X.-L. Fan and Q.-H. Zhang: Existence of solutions for $p(x)$ -Laplacian Dirichlet problem, *Nonlinear Anal.* **52** (2003), 1843–1852.
- [19] X.-L. Fan and D. Zhao: A class of De Giorgi type and Hölder continuity, *Nonlinear Anal.* **36** (1999), 295–318.
- [20] X.-L. Fan and D. Zhao: The quasi-minimizer of integral functionals with $m(x)$ growth conditions, *Nonlinear Anal.* **39** (2000), 807–816.
- [21] X.-L. Fan and D. Zhao: On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, *J. Math. Anal. Appl.* **263** (2001), 424–446.
- [22] X.-L. Fan, Y. Zhao and D. Zhao: Compact imbedding theorems with symmetry of Strauss-Lions type for the space $W^{1,p(x)}(\Omega)$, *J. Math. Anal. Appl.* **255** (2001), 333–348.
- [23] A. Fiorenza: A mean continuity type result for certain Sobolev spaces with variable exponent, *Commun. Contemp. Math.* **4** (2002), no. 3, 587–605.
- [24] T.C. Halsey: Electrorheological fluids, *Science* **258** (1992), 761–766.
- [25] P. Harjulehto and P. Hästö: A capacity approach to the Poincaré inequality and Sobolev imbedding in variable exponent Sobolev space, *Rev. Mat. Complut.*, to appear.
- [26] P. Harjulehto and P. Hästö: Lebesgue points in variable exponent spaces, preprint (2003).
- [27] P. Harjulehto, P. Hästö and M. Koskenoja: The Dirichlet energy integral on intervals in variable exponent Sobolev spaces, preprint (2003).
- [28] P. Harjulehto, P. Hästö, M. Koskenoja and S. Varonen: Sobolev capacity on the space $W^{1,p(\cdot)}(\mathbb{R}^n)$, *J. Function Spaces Appl.* **1** (2003), no. 1, 17–33.
- [29] P. Harjulehto, P. Hästö, M. Koskenoja and S. Varonen: The Dirichlet energy integral and variable exponent Sobolev spaces with zero boundary values, preprint (2003).
- [30] P. Harjulehto, P. Hästö, M. Koskenoja and S. Varonen: Variable Sobolev capacity and the bounds on the exponent, preprint (2003).
- [31] P. Harjulehto, P. Hästö and M. Pere: Variable exponent Lebesgue spaces on metric spaces: The Hardy–Littlewood maximal operator, preprint (2003).
- [32] J. Heinonen, T. Kilpeläinen and O. Martio: *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 1993.
- [33] H. Hudzik: On generalized Orlicz-Sobolev space, *Funct. Approximatio Comment. Math.* **4** (1976), 37–51.
- [34] H. Hudzik: A generalization of Sobolev spaces. I, *Funct. Approximatio Comment. Math.* **2** (1976), 67–73.
- [35] H. Hudzik: A generalization of Sobolev spaces. II, *Funct. Approximatio Comment. Math.* **3** (1976), 77–85.
- [36] V. Kokilashvili and S. Samko: Singular integrals in weighted Lebesgue spaces with variable exponent, *Georgian Math. J.* **10** (2003), no. 1, 145–156.
- [37] V. Kokilashvili and S. Samko: Singular integral equations in the Lebesgue spaces with variable exponent, *Proc. A. Razmadze Math. Inst.* **131** (2003), 61–78.

- [38] V. Kokilashvili and S. Samko: Singular integrals and potentials in some Banach function spaces with variable exponent, *J. Function Spaces Appl.* **1** (2003), no. 1, 45–59.
- [39] V. Kokilashvili and S. Samko: Maximal and fractional operators in weighted $L^{p(x)}$ spaces, *Rev. Mat. Iberoamericana*, to appear.
- [40] O. Kováčik and J. Rákosník: On spaces $L^{p(x)}$ and $W^{1,p(x)}$, *Czechoslovak Math. J.* **41(116)** (1991), 592–618.
- [41] P. Marcellini: Regularity and existence of solutions of elliptic equations with p, q -growth conditions, *J. Differential Equations* **50** (1991), no. 1, 1–30.
- [42] G. Mingione and D. Mucci: Integral functionals and the gap problem: sharp bounds for relaxation and energy concentration, preprint (2003).
- [43] J. Musielak: *Orlicz Spaces and Modular Spaces*, Springer-Verlag, Berlin, 1983.
- [44] A. Nekvinda: Hardy-Littlewood maximal operator on $L^{p(x)}(\mathbb{R}^n)$, *Math. Inequal. Appl.*, to appear.
- [45] C. Pfeiffer, C. Mavroidis, Y. Bar-Cohen, B. Dolgin: Electrorheological fluid based force feedback device, pp. 88–99 in *Proceedings of the 1999 SPIE Telemicromanipulator and Telepresence Technologies VI Conference* (Boston, MA), M. Stein (ed.), Vol. **3840** (1999).
- [46] L. Pick and M. Růžička: An example of a space $L^{p(x)}$ on which the Hardy-Littlewood maximal operator is not bounded, *Expo. Math.* **19** (2001), 369–371.
- [47] J. Rákosník: Sobolev inequality with variable exponent, pp. 220–228 in *Function Spaces, Differential Operators and Nonlinear Analysis* (Syöte, Finland, 1999), V. Mustonen and J. Rákosník (eds.), Mathematical Institute of the Academy of Sciences of the Czech Republic, Prague, 2000.
- [48] M. Růžička: *Electrorheological Fluids: Modeling and Mathematical Theory*, Springer-Verlag, Berlin, 2000.
- [49] S. Samko: Convolution type operators in $L^{p(x)}$, *Integr. Transform. and Special Funct.* **7** (1998), no. 1–2, 123–144.
- [50] S. Samko: Convolution and potential type operators in $L^{p(x)}(\mathbb{R}^n)$, *Integr. Transform. and Special Funct.* **7** (1998), no. 3–4, 261–284.
- [51] S. Samko: Denseness of $C_0^\infty(\mathbb{R}^n)$ in the generalized Sobolev spaces $W^{m,p(x)}(\mathbb{R}^n)$, pp. 333–342 in *Direct and Inverse Problems of Mathematical Physics (Newark, DE, 1997)*, Int. Soc. Anal. Appl. Comput. **5**, Kluwer Acad. Publ., Dordrecht, 2000.
- [52] I. I. Sharapudinov: On the topology of the space $L^{p(\cdot)}([0; 1])$, *Math. Notes* **26** (1979), no. 3–4, 796–806. [translation of *Mat. Zametki* **26** (1978), no. 4, 613–632.]
- [53] V. V. Zhikov: Averaging of functionals of the calculus of variations and elasticity theory, *Math. USSR-Izv.* **29** (1987), no. 1, 33–66. [translation of *Izv. Akad. Nauk SSSR Ser. Mat.* **50** (1986), no. 4, 675–710, 877.]

P. Harjulehto

DEPARTMENT OF MATHEMATICS
P.O. Box 4 (YLIOPISTONKATU 5)
FIN-00014 UNIVERSITY OF HELSINKI
FINLAND

P. Hästö

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MICHIGAN
ANN ARBOR, MI 48109-1109
USA

Research group home-page:

<http://www.math.helsinki.fi/analysis/varsobgroup>

E-mail addresses:

petteri.harjulehto@helsinki.fi
peter.hasto@helsinki.fi